



## Annex: Future climate projections for Tanzania

This annex provides the methods and extra figures to support the brief on future climate projections for Tanzania<sup>1</sup>, which provides an overview of future climate change for the country using the latest available climate model simulations. A two-page summary<sup>2</sup> is also available highlighting key findings.

### Data

#### Observed data

The observations used in the briefs included gridded rainfall and temperature data. We used the CHIRPS v2.0 (Climate Hazards group Infrared Precipitation with Stations) dataset<sup>3</sup> for daily rainfall at a resolution of ~5 km (0.05° x 0.05°) for the period 1981-2016. CHIRPS is a combination of satellite-based rainfall estimates and station observations and has been used for various analyses for Africa<sup>4,5,6</sup>. For temperature, we used the Climate Research Unit (CRU) TS v. 3.24.01<sup>7</sup> monthly data at a resolution of ~50 km (0.5° x 0.5°) for the period 1901-2015, which has been widely used for Africa<sup>8,9,10</sup>.

#### Historical and future climate data

We used the daily temperature and rainfall simulations for historical (1950-2005) and future (2006-2099) periods from 34 General Circulation Models (GCMs) listed in Table 1. These models are from the Coupled Model Intercomparison Project phase 5 (CMIP5) corresponding to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (AR5 IPCC). The historical simulations represent simulated climate variability from the mid-19th century to early 21st century, driven by anthropogenic and natural forcings. As the level of present emissions is just above the Representative Concentration Pathway (RCP) 8.5, we used climate simulations based on the high-emission business as usual pathway, RCP8.5<sup>11,12</sup>. Other lower emission RCPs are available in the CMIP5 database, but we only show results for RCP8.5 here.

#### Methods

Figure 1 shows the elevation and location map of Tanzania. For developing the climate brief, both temperature and rainfall variables have been extracted

over the domain – 0.25°S-12.25°S latitude and 28.75°E-40.75°E longitude for covering the geographical extent of Tanzania.

#### Downscaling of climate data

To analyse variations in future rainfall and temperature at a fine spatial resolution, we used the delta change method<sup>13,14</sup>. This method is widely used for downscaling coarse resolution GCM projections to derive information at a finer spatial scale<sup>15</sup> for climate change impact modelling (e.g. hydrological and crop modelling) studies, which require temperature and rainfall changes at finer spatial resolution<sup>16,17,18</sup>. In this method, a change factor is applied to the observed climatology of temperature and rainfall. This change factor represents the climate change signal as derived from the climate models, and is calculated as a difference of mean changes in the future and historical climate simulations of a GCM (Equation 1)<sup>19</sup>. The derived time series provides higher resolution information consistent with future projections of a changing climate<sup>20</sup>.

$$P_{\text{new}} = P_{\text{Obs}} + (P_{\text{futr}} - P_{\text{hist}}) \quad (1)$$

#### About FCFA

Future Climate for Africa (FCFA) aims to generate fundamentally new climate science focused on Africa, and to ensure that this science has an impact on human development across the continent.


[www.futureclimateafrica.org](http://www.futureclimateafrica.org)

where,  $P_{new}$ ,  $P_{obs}$ ,  $P_{futr}$  and  $P_{hist}$  represents the new time series, observations in the historical period, raw GCM output for the future period and raw GCM output for the historical period, respectively.  $P_{futr} - P_{hist}$  represents the change factor. Like other downscaling methods, there are advantages and limitations of using the delta change method. While it preserves the general climate change signal, it does not capture change in variance<sup>21,22</sup> or account for local climatic variation<sup>23,24</sup>. The method requires observations for the representative period<sup>25</sup>, which can be challenging in data scarce regions. For this study, the 30-year period from the historical simulations used in Equation 1 (1976-2005) is different from the available observations (1981-2010).

We present results for change in mean annual rainfall and temperature for all 34 CMIP5 models separately, and the Multi-Model Ensemble (MME) of historical and future simulations of temperature and rainfall from 34 CMIP5 models to show spatial patterns of change. We averaged monthly rainfall and temperature over the period

1976-2005 for historical simulations, medium-term (2021-2050), long-term (2070-2099) for future simulations and observed rainfall and temperature over the period 1981-2010. We obtained monthly change factors (12 each) for medium and long-term periods. The change factors were interpolated using bi-cubic interpolation and were added to CHIRPS and CRU observations for preparing new time series for medium and long-term future periods for rainfall and temperature, respectively. Using the monthly change factors we derived annual and seasonal change factors to address the seasonal climatic specificities of Tanzania.

For Tanzania, we used March to May (MAM) as the main rainy season known as the long rains, October to December (OND) known as the short rains and October through March in the following year (ONDJFM) to capture seasonality in central and western Tanzania. These seasons were considered important by the stakeholders, particularly the Ministry of Water and Irrigation and the Rufiji Water Basin Office in Tanzania.




**Table 1: List of 34 CMIP5 climate models used for analysing future climate change for Tanzania**

	Model	Modelling Centre/ Group
1	ACCESS1-0	Commonwealth Scientific and Industrial Research Organization (CSIRO) and Bureau of Meteorology (BOM), Australia
2	ACCESS1-3	
3	bcc-csm1-1	Beijing Climate Center, China Meteorological Administration, China
4	bcc-csm1-1-m	
5	BNU-ESM	College of Global Change and Earth System Science, Beijing Normal University, China
6	CanESM2	Canadian Centre for Climate Modelling and Analysis, Canada
7	CCSM4	University of Miami – RSMAS, United States
8	CESM1-BGC	Community Earth System Model Contributors, NSF–DOE–NCAR, United States
9	CESM1-CAM5	
10	CMCC-CESM	Centro Euro-Mediterraneo per I Cambiamenti Climatici, Italy
11	CMCC-CM	
12	CMCC-CMS	
13	CNRM-CM5	Centre National de Recherches Météorologiques, France
14	CSIRO-Mk3-6-0	CSIRO - Queensland Climate Change Centre of Excellence, Australia
15	EC-EARTH	Irish Centre for High-End Computing (ICHEC), Ireland
16	FGOALS-g2	LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences and CESS, Tsinghua University, China
17	GFDL-CM3	NOAA Geophysical Fluid Dynamics Laboratory, United States
18	GFDL-ESM2G	
19	GFDL-ESM2M	
20	HadGEM2-AO	National Institute of Meteorological Research/Korea Meteorological Administration, South Korea
21	HadGEM2-CC	Met Office Hadley Centre, United Kingdom
22	HadGEM2-ES	
23	INM-CM4	Institute for Numerical Mathematics, Russia
24	IPSL-CM5A-LR	Institut Pierre-Simon Laplace, France
25	IPSL-CM5A-MR	
26	IPSL-CM5B-LR	
27	MIROC5	Japan Agency for Marine-Earth Science and Technology, Atmosphere and Ocean Research Institute (The University of Tokyo), and National Institute for Environmental Studies, Japan
28	MIROC-ESM	
29	MIROC-ESM-CHEM	
30	MPI-ESM-LR	Max Planck Institute for Meteorology, Germany
31	MPI-ESM-MR	
32	MRI-CGCM3	Meteorological Research Institute, Japan
33	MRI-ESM1	
34	NorESM1-M	Norwegian Climate Centre, Norway


## Recent climate variability and extremes from observations

Figure 2: CHIRPS observed annual and seasonal rainfall trend (linear trend by grid cell in mm/year) for 1981-2016. Seasons are March to May (MAM), October to December (OND), and October to the following March (ONDJFM)


The outline of the Rufiji river basin is shown in black.



**Figure 3: CHIRPS observed mean seasonal precipitation anomaly (%) for the driest year 2005, wettest year 2006/07 and recent dry year 2014/15 with respect to the mean for 1981-2016 MAM and ONDJFM seasons**



**Figure 4: CRU observed annual and seasonal mean temperature (°C) for 1976-2005. Seasons are March to May (MAM), October to December (OND), and October to the following March (ONDJFM)**




## Projections of future climate


We use the daily temperature and rainfall simulations for historical (1950-2005) and future (2006-2099) periods from 34 GCMs from CMIP5 corresponding to the IPCC's AR5. We present climate simulations based only on a high greenhouse gas emission pathway, RCP8.5 (projections with other rates of emissions are available).

## Rainfall

**Figure 5: Per cent change in annual mean rainfall for all Tanzania between the GCM simulated current period (1976-2005) and 2070-2099 for 34 GCMs**



**Figure 6: Mean annual and seasonal precipitation change (%) for the 2090s (2070-2099) compared to current period 1976-2005 using ensemble mean 34 CMIP5 models for annual, OND, ONDJFM and MAM**



## Temperature

Figure 7: Change in annual mean temperature (°C) for all Tanzania between the GCM simulated current period (1976-2005) and 2021-2050 for 34 GCMs







Figure 8: Change in annual mean temperature (°C) between the GCM simulated current period (1976-2005) and 2070-99 for 34 GCMs




**Figure 9: Mean seasonal temperature change (°C) for near-term 2021-2050 compared to current period 1976-2005 using ensemble mean of 34 CMIP5 models for annual, OND, ONDJFM and MAM**



**Figure 10: Mean seasonal temperature change (°C) for the long-term 2070-2099 compared to current period 1976-2005 using ensemble mean of 34 CMIP5 models for annual, OND, ONDJFM and MAM**



**Figure 11: Time series of mean annual temperature (°C) for 34 CMIP5 models and their ensemble (bold red line) for the period 1950-2099 and CRU observations (bold black line) for the period 1950-2014**



## Endnotes

- 1 FCFA (2017a) Country climate brief. *Future climate projections for Tanzania: Cape Town: Future Climate for Africa*. www.futureclimateafrica.org/resource/future-climate-projections-for-tanzania/
- 2 FCFA (2017b) Country climate brief. *Summary: Future climate projections for Tanzania: Cape Town: Future Climate for Africa*. www.futureclimateafrica.org/resource/future-climate-projections-for-tanzania/
- 3 Funk, C., Peterson, P., Landsfeld, M., Pedreros, D., Verdin, J., Shukla, S., Husak, G., Rowland, J., Harrison, L., Hoell, A. and Michaelsen, J., (2015) The climate hazards infrared precipitation with stations--a new environmental record for monitoring extremes. *Scientific Data*, 2, p. 150066.
- 4 Badr, H. S., Dezfuli, A. K., Zaitchik, B. F., & Peters-Lidard, C. D. (2016) Regionalizing Africa: Patterns of Precipitation Variability in Observations and Global Climate Models. *Journal of Climate*, 29(24), p. 027-9043.
- 5 McNally, A., Arsenault, K., Kumar, S., Shukla, S., Peterson, P., Wang, S., Funk, C., Peters-Lidard, C.D. & Verdin, J. P. (2017) A land data assimilation system for sub-Saharan Africa food and water security applications. *Scientific Data*, 4, p. 170012.
- 6 Sossa, A., Liebmann, B., Bladé, I., Al-lured, D., Hendon, H. H., Peterson, P., & Hoell, A. (2017) Statistical Connection between the Madden-Julian Oscillation and Large Daily Precipitation Events in West Africa. *Journal of Climate*, 30(6), p. 1999-2010.
- 7 Harris, I.P.D.J., Jones, P.D., Osborn, T.J. and Lister, D.H., (2014) Updated high resolution grids of monthly climatic observations--the CRU TS3. 10 Dataset. *International Journal of Climatology*, 34(3), p. 623-642.
- 8 Fotso-Nguemo, T. C., Vondou, D. A., Tchawoua, C., & Haensler, A. (2016) Assessment of simulated rainfall and temperature from the regional climate model REMO and future changes over Central Africa. *Climate Dynamics*, p. 1-21.
- 9 Camberlin, P. (2017) Temperature trends and variability in the Greater Horn of Africa: interactions with precipitation. *Climate Dynamics*, 48(1-2), p. 477-498.
- 10 Sossa, A. et al. (2017) Op. cit.
- 11 Riahi, K., Rao, S., Krey, V., Cho, C., Chirkov, V., Fischer, G., Kindermann, G., Nakicenovic, N. & Rafaj, P. (2011) RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Climatic Change*, 109(1-2), p. 33.
- 12 Sanford, T., Frumhoff, P. C., Luers, A., & Gullede, J. (2014) The climate policy narrative for a dangerously warming world. *Nature Climate Change*, 4(3), p. 164-166.
- 13 Tabor, K., & Williams, J. W. (2010) Globally downscaled climate projections for assessing the conservation impacts of climate change. *Ecological Applications*, 20(2), p. 554-565.
- 14 Rätty, O., Räisänen, J., & Ylhäisi, J. S. (2014) Evaluation of delta change and bias correction methods for future daily precipitation: intermodel cross-validation using ENSEMBLES simulations. *Climate Dynamics*, 42(9-10), p. 2287-2303.
- 15 Keller, D. E., Fischer, A. M., Liniger, M. A., Appenzeller, C., & Knutti, R. (2017) Testing a weather generator for downscaling climate change projections over Switzerland. *International Journal of Climatology*, 37, p. 928-942.
- 16 Teutschbein, C., Wetterhall, F., & Seibert, J. (2011) Evaluation of different downscaling techniques for hydrological climate-change impact studies at the catchment scale. *Climate Dynamics*, 37(9-10), p. 2087-2105.
- 17 Hawkins, E., Osborne, T. M., Ho, C. K., & Challinor, A. J. (2013) Calibration and bias correction of climate projections for crop modelling: an idealised case study over Europe. *Agricultural and Forest Meteorology*, 170, p. 19-31.
- 18 House, A. R., Thompson, J. R., & Acreman, M. C. (2016) Projecting impacts of climate change on hydrological conditions and biotic responses in a chalk valley riparian wetland. *Journal of Hydrology*, 534, p. 178-192.
- 19 Tabor, K., & Williams, J. W. (2010) Op. cit.
- 20 Rätty, O. et al. (2014) Op. cit.
- 21 Teutschbein, C., Wetterhall, F., & Seibert, J. (2011) Op. cit.
- 22 Addor, N., Rössler, O., Köplin, N., Huss, M., Weingartner, R., & Seibert, J. (2014) Robust changes and sources of uncertainty in the projected hydrological regimes of Swiss catchments. *Water Resources Research*, 50(10), p. 7541-7562.
- 23 Onyutha, C., Tabari, H., Rutkowska, A., Nyeko-Ogiramoi, P., & Willems, P. (2016). Comparison of different statistical downscaling methods for climate change rainfall projections over the Lake Victoria basin considering CMIP3 and CMIP5. *Journal of Hydro-environment Research*, 12, p. 31-45.
- 24 Keller, D. E. et al. (2017) Op. cit.
- 25 Hawkins, E. et al. (2017) Op. cit.
- 26 DIVA GIS www.diva-gis.org/gdata

---

## About Future Climate for Africa

Future Climate for Africa (FCFA) aims to generate fundamentally new climate science focused on Africa, and to ensure that this science has an impact on human development across the continent. This brief was written by Neha Mittal of the UMFULA research team. You can find out more about their work under 'research teams' on [www.futureclimateafrica.org](http://www.futureclimateafrica.org).



Funded by:



[www.futureclimateafrica.org](http://www.futureclimateafrica.org)

e: [info@futureclimateafrica.org](mailto:info@futureclimateafrica.org)

t: +2721 4470211

---

This document is an output from a project funded by the UK Department for International Development (DFID) and the Natural Environment Research Council (NERC) for the benefit of developing countries and the advance of scientific research. However, the views expressed and information contained in it are not necessarily those of, or endorsed by DFID or NERC, which can accept no responsibility for such views or information or for any reliance placed on them. This publication has been prepared for general guidance on matters of interest only, and does not constitute professional advice. You should not act upon the information contained in this publication without obtaining specific professional advice. No representation or warranty (expressed or implied) is given as to the accuracy or completeness of the information contained in this publication, and, to the extent permitted by law, the Climate and Development Knowledge Network's members, the UK Department for International Development ('DFID'), the Natural Environment Research Council ('NERC'), their advisors and the authors and distributors of this publication do not accept or assume any liability, responsibility or duty of care for any consequences of you or anyone else acting, or refraining to act, in reliance on the information contained in this publication or for any decision based on it.

Copyright © 2017, Climate and Development Knowledge Network. All rights reserved.